Bold type indicates authors of letters, commentaries, editorials, and Resource Corners; (No.) indicates issue number.

The ACTIVE Writing Group on behalf of the ACTIVE Investigators, 58
Acuin JM, 48
Ahmed A, 78
Algra A, 57
Alonso-Coello P, 1
Alper BS, A-10 (No. 3)
Anderson GL, 4
Antman EM, 30, 52
Aragona G, 49
Arden NK, 15
Arnold MJ, 2
Assanelli E, 63
Aujesky D, 53
Aumann JL, 39
Avorn J, 80

Baigent C, 66
Baime MJ, 17
Ballew KA, 33
Barnett AH, 67
Barnsley L, 15
Barratt A, A-10 (No. 1)
Barrett-Connor E, 73
Basu A, 67
Bates ER, 9, 59
Bauer DC, 71
Beaupre LA, 10
Beck SJ, 40
Bent S, 12
Beresford SA, 6
Berger PB, 61
Bernard GR, 69, 70
Berry E, 77
Best JJ, 77
Bhatia RS, 78
Bhatt DL, 33
Blaivas AJ, 43
Boissel JP, 64
Bønaa KH, 3
Bongartz T, 65
Borzak S, 29
Boudville N, 79
Braunwald E, 52
Briel M, 61
Brochard L, 70
Brok J, 47
Busto UE, 14

Caan B, 6
Carey BC, 61
Carey T, 74
Carusone SC, 81
Casas JP, 36
Casazza G, 21
Chalmers I, A-8–A-9 (No. 1)
Chan FK, 49
Chang TT, 18
Chappell FM, 77
Chatellier G, 23
Chen W, 47
Cheng EY, 22
Cherkin DC, 16
Chlebowski RT, 6
Chrolavicius S, 29, 31
Chua W, 36
Clarfield AM, 14
Clarkson JE, 75
Clase C, 36
Clegg DO, 17
Coates AS, 11
Colli A, 21
Collins P, 73
Connolly S, 58
Cook D, 68
Cornell JE, 37, 38
Costantino JP, 72
Craig JC, 35
Craig MC, 35
Cranney A, 71
Crockett S, 44
Cucherat M, 32

Dal Bó N, 49
Danchin N, 32
Davenport C, 22
de Man R, 18
d'Emden M, 62
Demicheli V, 45
Dew MA, 13
Deyo RA, 16
Di Mario F, 49
Dixon JB, 41
Doucette S, 24
Doust J, 22
Dreyer M, 67
Duncan DG, 40
Dunn A, 23, 25
Duque G, 40

Eden OB, 75
Eikelboom JW, 9
Ellison III RT, 46
Elmer PJ, 42
Ernst E, 20
Erro J, 16
Eurich DT, 80
Evans AT, 16

Fay S, 50
Fergusson D, 24
Fleetham J, 43
Fletcher RH, 4
Fouque D, 64
Fowler SE, 76
Fox KA, 33
Fraquelli M, 21
Frederick PD, 52
Furlan AD, 74

Gamble C, 48
Garcia-Alamino JM, 1
Garg AX, 36
Gass M, 4
Gerson LB, 44
Giannini EG, 21
Gill SS, 10
Gish RG, 18
Ghosh AK, A-11 (No. 2)
Glass J, 14
Gluud C, 47
Gluud LL, 47
Godwin J, 66
Goeree R, 81
Goodman LR, 76
Gould MK, 53
Green ML, A-8–A-10 (No. 2)
Guitteau M, 42

Hacke W, 33
Hadler NM, 16
Hak E, 68
Halkes PH, 57
Harbord R, 51
Harris CL, 17
Hart R, 58
Haynes RB, A-8–A-9 (No. 3)
Heneghan C, A-11 (No.1), 1
Hermans G, 34
Herrmann N, 14
Hirsch C, 50
Hochman JS, 59
Holloway RG, 77
Hood K, 40
Hou FF, 19
Howard BV, 6
Hsia J, 6
Hsu C, 19, 79
Hudson MP, 8

Jackson RD, 4
James M, 63
Janssens E, 39
Jefferson T, 45
Johansen A, 40
Johnson KC, 6
Jones CA, 10

Kahn SR, 20
Kaltenbach T, 44
Kane C, 12
Kappelle LJ, 57
Kastrati A, 8
Kearney PM, 66
Kelly AG, 77
Keshaviah A, 11
Khan N, 60
Knoll G, 79
Knopp RH, 62
Koeman M, 68
Kotchen JM, 4
Koudstaal PJ, 57
Krahn M, 47
Krebs E, 74
Kudva YC, 67
Kwok YT, 22

LaCroix AZ, 4
Lanctôt KL, 14
Lange P, 67
Langley JM, 45
Lasserson TJ, 43
Laupacis A, 66
Laurie C, 41
Laville M, 64
Lawrence VA, 37, 38
Le Gal G, 25
Le Manach CP, 53
Lee DS, 78
Lim J, 43
Loeb M, 81
Lonn E, 2
Loukogeorgakis S, 36
Lous J, 48

Macfadyen CA, 48
Mailis-Gagnon A, 74
Main C, 51
Majumdar SR, 80
Manns B, 63
Marana I, 63
Marenzi G, 63
Massel D, 30–31
McAlister FA, 60
McCabe CH, 30
McFarland LV, 46
McFarlane PA, 35
Mehilli J, 8
Mehta SR, 9, 29, 31
Miglioretti DL, 16
Montori VM, 62
Morrow DA, 30, 52
Mosca L, 73

Neumann FJ, 8
Njølstad I, 3

Obarzanek E, 42
O'Brien PE, 41
Owen C, 24

Padwal R, 41
Papaioannou A, 71
Perrier A, 24, 76
Pittler MH, 20
Pocock SJ, 62
Pogue J, 58
Pollock BG, 13
Prasad R, 79
Prentice RL, 6
Price C, 15
Prowell TM, 11

Quinlan DJ, 9
Qushmaq I, 69

Rabbat CG, 64
Reading I, 15
Reda DJ, 17
Reichen J, 18
Reynolds CF 3rd, 13
Righini M, 25
Ritenbaugh C, 6
Rivetti D, 45
Rockwood K, 50
Rosenberg M, 42
Roy PM, 25, 53
Ruest A, 68

Sacks HS, 81
Sandoval JA, 74
Saunders LD, 10
Schena FP, 35
Schoenfeld D, 70
Schünemann HJ, 39
Schwartz GG, 61
Scott IA, 52
Serdarevic-Pehar M, 67
Sharma P, 44
Shaw JE, 75
Shekelle P, 2–3
Sherman KJ, 16
Shinohara K, 12
Shoor S, 65
Simpson SH, 80
Sleeper LA, 59
Sloan MA, 58
Smetana GW, 37, 38
Smilde JG, 62
Song X, 50
Spandorfer J, 1
Sproule BA, 14
Stearns V, 11, 72
Stein PD, 76
Steinberg DI, 37–38
Stone PH, 32
Strippoli GF, 35
Sutton AJ, 65
Sweeting MJ, 65
Thacker HL, 6, 73
Thompson BT, 34, 70
Thompson PL, 61
Thuillez C, 32
Thürlimann B, 11
Tirschwell D, 57
Touzé E, 23
Tran H, 24
Tu JV, 78
Tunks E, 74

Ueland PM, 3

Van den Berghe G, 34
van der Ven AJ, 68
van Gijn J, 57
Van Horn L, 6
van Noord JA, 39
Varenne O, 23
Vinson DC, A-10 (No. 3)
Vogel VG, 72
Vollmer WM, 42

Wactawski-Wende J, 4
Wani S, 44
Wardlaw JM, 77
Wartolowska K, 77
Webb JG, 59
Weinshenker BG, 51
Weiss R, 60
Wells PS, 24
Wheeler AP, 69, 70
Whiting P, 51
Wickerham DL, 72
Wiedemann HP, 69, 70
Williams Jr JW, 13
Wilmer A, 34
Wilt TJ, 12
Wiviott SD, 52
Wong VW, 49
Worthington HV, 75
Wright J, 43

Yusuf S, 2, 29, 31

Zhang GH, 19
Zhang X, 19
Zytaruk N, 71

A

Acetylcysteine
- N-acetylcysteine prevented contrast-medium–induced nephropathy in primary angioplasty, 63

Adrenergic beta-agonists
- Formoterol in addition to tiotropium improved airflow obstruction in chronic obstructive pulmonary disease, 39

Adrenergic beta-antagonists
- Review: β-blockers differ in their efficacy for preventing major cardiovascular events in younger and older patients, 60

Aesculus
- Review: Horse chestnut seed extract is effective for symptoms of chronic venous insufficiency, 20

Albuminuria
- Review: ACE inhibitors delay onset of microalbuminuria in diabetes without nephropathy and reduce mortality in diabetic nephropathy, 35

Alpha-fetoproteins
- Review: MRI is more sensitive but less specific than ultrasonography or spiral CT for diagnosis of hepatocellular carcinoma, 21

Alzheimer disease
- Galantamine was effective in mild-to-moderate Alzheimer disease, 50

Analgesics, opioid
- Review: Opioids are more effective than placebo but not other analgesics for chronic noncancer pain, 74

Androgen antagonists
- Saw palmetto did not differ from placebo for benign prostatic hyperplasia in men, 12

Angina
- Abciximab reduced death, MI, and urgent target vessel revascularization in non–ST-segment elevation acute coronary syndromes, 8
- Fondaparinux was noninferior to enoxaparin for death, MI, and refractory ischemia but reduced bleeding in angina and non-STEMI, 31
- Review: Early statin therapy does not reduce the composite endpoint of death, MI, or stroke in acute coronary syndromes, 61

Angiotensin-converting enzyme inhibitors
- Benazepril was effective and safe for advanced chronic kidney disease without diabetes, 19
- Review: ACE inhibitors reduce mortality and cardiovascular endpoints in stable coronary artery disease, 32
- Review: ACE inhibitors delay onset of microalbuminuria in diabetes without nephropathy and reduce mortality in diabetic nephropathy, 35
- Review: The renoprotective effects of ACE inhibitors and ARBs independent of blood pressure control are uncertain, 36

Anti-bacterial agents
- Bovine lactoferrin added to triple therapy increased *Helicobacter pylori* eradication rate, 49
- Review: Probiotics are effective for prevention of antibiotic-associated diarrhea and treatment of *Clostridium difficile* disease, 46
- Review: Topical quinolones are better than systemic antibiotics for chronic suppurative otitis media at up to 2 weeks, 48

Antibodies, monoclonal
- Abciximab reduced death, MI, and urgent target vessel revascularization in non–ST-segment elevation acute coronary syndromes, 8
- Review: Anti–tumor necrosis factor antibody therapy for rheumatoid arthritis increases risk for serious infection and malignancy, 65

Anticoagulants
- Clopidogrel plus aspirin was inferior to oral anticoagulation for preventing vascular events in atrial fibrillation, 58
- Enoxaparin for 7 days was better than unfractionated heparin for 2 days for reducing death and MI but not bleeding in STEMI, 30
- Fondaparinux reduced death or reinfarction in acute ST-segment elevation myocardial infarction, 29
- Fondaparinux was noninferior to enoxaparin for death, MI, and refractory ischemia but reduced bleeding in angina and non-STEMI, 31
- Review: Self-monitoring increases the efficacy and safety of anticoagulant therapy, 1

Antifungal agents
- Review: Prophylactic interventions reduce oral mucositis in patients with cancer receiving chemotherapy or radiation therapy, 75

Antihypertensive agents
- Review: ACE inhibitors delay onset of microalbuminuria in diabetes without nephropathy and reduce mortality in diabetic nephropathy, 35
- Review: The renoprotective effects of ACE inhibitors and ARBs independent of blood pressure control are uncertain, 36

Antiinfective agents
- Oral decontamination with chlorhexidine reduced ventilator-associated pneumonia in high-risk patients, 68

Anti-inflammatory agents, non-steroidal
- Aspirin plus dipyridamole was more effective than aspirin alone for preventing vascular events after minor cerebral ischemia, 57
- Review: Selective COX-2 inhibitors increase vascular events more than placebo and naproxen, but not more than other NSAIDs, 66

Antineoplastic agents
- Disease-free survival was greater with letrozole than tamoxifen in postmenopausal women with early breast cancer, 11
- Review: Prophylactic interventions reduce oral mucositis in patients with cancer receiving chemotherapy or radiation therapy, 75

Antirheumatic agents
- Review: Anti–tumor necrosis factor antibody therapy for rheumatoid arthritis increases risk for serious infection and malignancy, 65

Antiviral agents
- Entecavir was more effective than lamivudine in HbeAg-positive chronic hepatitis B, 18
- Review: Antiviral agents reduce risk for influenza in healthy adults and alleviate symptoms faster than placebo, 45
- Review: Ribavirin is not better than placebo in chronic hepatitis C infection, 47

Arthritis, rheumatoid
- Review: Anti–tumor necrosis factor antibody therapy for rheumatoid arthritis increases risk for serious infection and malignancy, 65

Aspirin
- Aspirin plus dipyridamole was more effective than aspirin alone for preventing vascular events after minor cerebral ischemia, 57
- Clopidogrel plus aspirin did not differ from aspirin alone for reducing MI, stroke, and CV death in high-risk atherothrombosis, 33
- Clopidogrel plus aspirin was inferior to oral anticoagulation for preventing vascular events in atrial fibrillation, 58

Atherosclerosis
- Review: Selective COX-2 inhibitors increase vascular events more than placebo and naproxen, but not more than other NSAIDs, 66

Atrial fibrillation
- Clopidogrel plus aspirin was inferior to oral anticoagulation for preventing vascular events in atrial fibrillation, 58

B

Benzazepines
Benazepril was effective and safe for advanced chronic kidney disease without diabetes, 19
Breast neoplasms
A low-fat dietary pattern intervention did not reduce incidence of breast cancer, colorectal cancer, or CVD in postmenopausal women, 6
Disease-free survival was greater with letrozole than tamoxifen in postmenopausal women with early breast cancer, 11
Raloxifene and tamoxifen had similar efficacy for preventing invasive breast cancer in women at increased risk, 72
Raloxifene produced both harms and benefits in postmenopausal women, with no reduction in cardiovascular disease risk, 73
Bronchodilator agents
Formoterol in addition to tiotropium improved airflow obstruction in chronic obstructive pulmonary disease, 39

C

Calcium carbonate
Calcium plus vitamin D did not prevent hip fracture or colorectal cancer in postmenopausal women, 4
Calcium channel blockers
Review: ACE inhibitors delay onset of microalbuminuria in diabetes without nephropathy and reduce mortality in diabetic nephropathy, 35
Candidiasis, oral
Review: Prophylactic interventions reduce oral mucositis in patients with cancer receiving chemotherapy or radiation therapy, 75
Carcinoma, hepatocellular
Review: MRI is more sensitive but less specific than ultrasonography or spiral CT for diagnosis of hepatocellular carcinoma, 21
Cardiovascular diseases
Clopidogrel plus aspirin did not differ from aspirin alone for reducing MI, stroke, and CV death in high-risk atherothrombosis, 33
Lowering homocysteine with folic acid and B vitamins did not prevent vascular events in vascular disease, 2
Carotid stenosis
Review: Noninvasive imaging techniques may be useful for diagnosing 70% to 99% carotid stenosis in symptomatic patients, 77
Catheterization
Pulmonary artery catheters and central venous catheters did not differ for mortality in acute lung injury, 70
Cerebrovascular accident
Aspirin plus dipyridamole was more effective than aspirin alone for preventing vascular events after minor cerebral ischemia, 57
Atorvastatin did not prevent cardiovascular events in type 2 diabetes, 62
Clopidogrel plus aspirin was inferior to oral anticoagulation for preventing vascular events in atrial fibrillation, 58
A low-fat dietary pattern intervention did not reduce incidence of breast cancer, colorectal cancer, or CVD in postmenopausal women, 6
Review: Early statin therapy does not reduce the composite endpoint of death, MI, or stroke in acute coronary syndromes, 61
Risks for myocardial infarction or nonstroke vascular death after ischemic stroke or transient ischemic attack were each 2% per year, 23
Chlorhexidine
Oral decontamination with chlorhexidine reduced ventilator-associated pneumonia in high-risk patients, 68
Clarithromycin
Bovine lactoferrin added to triple therapy increased *Helicobacter pylori* eradication rate, 49
Clostridium difficile
Review: Probiotics are effective for prevention of antibiotic-associated diarrhea and treatment of *Clostridium difficile* disease, 46
Colorectal neoplasms
Calcium plus vitamin D did not prevent hip fracture or colorectal cancer in postmenopausal women, 4
A low-fat dietary pattern intervention did not reduce incidence of breast cancer, colorectal cancer, or CVD in postmenopausal women, 6
Coronary disease
A low-fat dietary pattern intervention did not reduce incidence of breast cancer, colorectal cancer, or CVD in postmenopausal women, 6
Raloxifene produced both harms and benefits in postmenopausal women, with no reduction in cardiovascular disease risk, 73
Review: ACE inhibitors reduce mortality and cardiovascular endpoints in stable coronary artery disease, 32
Critical pathways
A nurse-led clinical pathway reduced hospitalizations in nursing home residents with pneumonia, 81
Cyclooxygenase 2 inhibitors
Review: Selective COX-2 inhibitors increase vascular events more than placebo and naproxen, but not more than other NSAIDs, 66

D

Depressive disorder, major
Maintenance treatment with paroxetine, but not psychotherapy, prevented recurrent major depression in older persons, 13
Diabetes mellitus
Adjunctive inhaled insulin before meals improved glycemic control more than adjunctive metformin in type 2 diabetes mellitus, 67
Atorvastatin did not prevent cardiovascular events in type 2 diabetes, 62
Lowering homocysteine with folic acid and B vitamins did not prevent vascular events in vascular disease, 2
Diabetic nephropathies
Review: ACE inhibitors delay onset of microalbuminuria in diabetes without nephropathy and reduce mortality in diabetic nephropathy, 35
Review: The renoprotective effects of ACE inhibitors and ARBs independent of blood pressure control are uncertain, 36
Diarrhea
Review: Probiotics are effective for prevention of antibiotic-associated diarrhea and treatment of *Clostridium difficile* disease, 46
Diet
A low-fat dietary pattern intervention did not reduce incidence of breast cancer, colorectal cancer, or CVD in postmenopausal women, 6
Review: A low-protein diet delays end-stage renal disease or death in stage 3 to 5 chronic kidney disease,064
Dietary supplements
Dietetic assistants improved postoperative clinical outcomes in older women with hip fracture, 40
Dipyridamole
Aspirin plus dipyridamole was more effective than aspirin alone for preventing vascular events after minor cerebral ischemia, 57

E

Education
Evaluating evidence-based practice performance, 31
Electrocardiography
Review: Electrocardiography, BNP, and N terminal-pro BNP are more sensitive than specific for chronic left ventricular systolic dysfunction, 22
The Thrombolysis In Myocardial Infarction risk index predicted in-hospital mortality in non–ST-segment elevation myocardial infarction, 52
Enoxaparin
Enoxaparin for 7 days was better than unfractionated heparin for 2 days for reducing death and MI but not bleeding in STEMI, 30
Fondaparinux was noninferior to enoxaparin for death, MI, and refractory ischemia but reduced bleeding in angina and non-STEMI, 31
Evidence-based medicine
Evaluating evidence-based practice performance (Editorial), A-8 (No. 2)
Evaluating evidence-based practice performance (Letter), A-10 (No. 3)
Clinical Epidemiology: How to do Clinical Practice Research (Resource Corner), A-11 (No. 1)
Evidence-based Medicine Toolkit (Resource Corner), A-11 (No. 2)
Of studies, syntheses, synopses, summaries, and systems: the "5S" evolution of information services for evidence-based health care decisions, 30
Selling Sickness (Resource Corner), A-10 (No. 1)
Why fair tests are needed: a brief history (Editorial), A-8 (No. 1)
Exercise therapy
Yoga improved function and reduced symptoms of chronic low-back pain more than a self-care book, 16

F

Fibrin fibrinogen degradation products
Review: The Wells clinical prediction guide and D-dimer testing predict deep venous thrombosis, 24
Fibrinolytic agents
Enoxaparin for 7 days was better than unfractionated heparin for 2 days for reducing death and MI but not bleeding in STEMI, 30
Review: Low-molecular-weight heparin is more effective than unfractionated heparin for preventing reinfarction in STEMI, 9
Fluid therapy
A conservative fluid management strategy did not affect risk for death but shortened duration of ventilation in acute lung injury, 69
Folic acid
Lowering homocysteine with folic acid and B vitamins did not prevent vascular events in vascular disease, 2
Lowering homocysteine with folic acid and B vitamins did not prevent vascular events after myocardial infarction, 3

G

Galantamine
Galantamine was effective in mild-to-moderate Alzheimer disease, 50
Gastroesophageal reflux
Review: Sparse evidence supports lifestyle modifications for reducing symptoms of gastroesophageal reflux disease, 44
Glucocorticoids
Lumbar epidural corticosteroid injections provided only short-term relief of symptoms and pain in unilateral sciatica, 15
Glucosamine
Glucosamine and chondroitin sulfate did not improve pain in osteoarthritis of the knee, 17
Guanine
Entecavir was more effective than lamivudine in HbeAg-positive chronic hepatitis B, 18

H

Heart failure, congestive
1-year mortality after first hospitalization for heart failure was similar in patients with preserved or reduced ejection fraction, 78
Helicobacter infections
Bovine lactoferrin added to triple therapy increased *Helicobacter pylori* eradication rate, 49
Heparin
Enoxaparin for 7 days was better than unfractionated heparin for 2 days for reducing death and MI but not bleeding in STEMI, 30
Review: Self-monitoring increases the efficacy and safety of anticoagulant therapy, 1
Hepatitis
Entecavir was more effective than lamivudine in HbeAg-positive chronic hepatitis B, 18
Review: Ribavirin is not better than placebo in chronic hepatitis C infection, 47
Hip fractures
Dietetic assistants improved postoperative clinical outcomes in older women with hip fracture, 40
Review: Strong evidence supports perioperative practices to reduce complications from hip fracture, 10
Hospital mortality
The Thrombolysis In Myocardial Infarction risk index predicted in-hospital mortality in non–ST-segment elevation myocardial infarction, 52
Hyperhomocysteinemia
Lowering homocysteine with folic acid and B vitamins did not prevent vascular events in vascular disease, 2
Lowering homocysteine with folic acid and B vitamins did not prevent vascular events after myocardial infarction, 3
Hypertension
Lifestyle recommendations plus the DASH diet reduced hypertension in patients with above-optimal blood pressure, 42
Review: β-blockers differ in their efficacy for preventing major cardiovascular events in younger and older patients, 60
Review: Healthy kidney donors may have a long-term increase in blood pressure beyond that associated with normal aging, 79
Hypnotics and sedatives
Review: Sedative–hypnotics increase adverse effects more than they improve sleep quality in older persons with insomnia, 14
Hypoglycemic agents
Intensive insulin therapy reduced morbidity but not mortality in patients in the medical intensive care unit, 34

I

Influenza, human
Review: Antiviral agents reduce risk for influenza in healthy adults and alleviate symptoms faster than placebo, 45
Information systems
Of studies, syntheses, synopses, summaries, and systems: the "5S" evolution of information services for evidence-based health care decisions, 30

Insulin
Adjunctive inhaled insulin before meals improved glycemic control more than adjunctive metformin in type 2 diabetes mellitus, 67
Intensive insulin therapy reduced morbidity but not mortality in patients in the medical intensive care unit, 34
Ischemic attack, transient
Risks for myocardial infarction or nonstroke vascular death after ischemic stroke or transient ischemic attack were each 2% per year, 23

K

Kidney
N-acetylcysteine prevented contrast-medium–induced nephropathy in primary angioplasty, 63
Review: Healthy kidney donors may have a long-term increase in blood pressure beyond that associated with normal aging, 79
Review: A low-protein diet delays end-stage renal disease or death in stage 3 to 5 chronic kidney disease, 64
Review: The renoprotective effects of ACE inhibitors and ARBs independent of blood pressure control are uncertain, 36

L

Lactoferrin
Bovine lactoferrin added to triple therapy increased *Helicobacter pylori* eradication rate, 49
Lamivudine
Entecavir was more effective than lamivudine in HbeAg-positive chronic hepatitis B, 18
Liver neoplasms
Review: MRI is more sensitive but less specific than ultrasonography or spiral CT for diagnosis of hepatocellular carcinoma, 21
Low back pain
Yoga improved function and reduced symptoms of chronic low-back pain more than a self-care book, 16
Lung diseases
Review: Lung expansion modalities and selective nasogastric decompression prevent postoperative pulmonary complications, 38
Review: Surgical site, advanced age, and comorbid conditions increase risk for postoperative pulmonary complications, 37

M

Magnetic resonance imaging
Review: Magnetic resonance imaging alone is of limited usefulness in diagnosing multiple sclerosis, 51
Review: MRI is more sensitive but less specific than ultrasonography or spiral CT for diagnosis of hepatocellular carcinoma, 21
Metformin
Adjunctive inhaled insulin before meals improved glycemic control more than adjunctive metformin in type 2 diabetes mellitus, 67
Mouth
Oral decontamination with chlorhexidine reduced ventilator-associated pneumonia in high-risk patients, 68
Multiple sclerosis
Review: Magnetic resonance imaging alone is of limited usefulness in diagnosing multiple sclerosis, 51
Myocardial infarction
Atorvastatin did not prevent cardiovascular events in type 2 diabetes, 62
Enoxaparin for 7 days was better than unfractionated heparin for 2 days for reducing death and MI but not bleeding in STEMI, 30
Fondaparinux reduced death or reinfarction in acute ST-segment elevation myocardial infarction, 29
Fondaparinux was noninferior to enoxaparin for death, MI, and refractory ischemia but reduced bleeding in angina and non-STEMI, 31
Lowering homocysteine with folic acid and B vitamins did not prevent vascular events after myocardial infarction, 3
Review: Early statin therapy does not reduce the composite endpoint of death, MI, or stroke in acute coronary syndromes, 61
Review: Low-molecular-weight heparin is more effective than unfractionated heparin for preventing reinfarction in STEMI, 9
Risks for myocardial infarction or nonstroke vascular death after ischemic stroke or transient ischemic attack were each 2% per year, 23
The Thrombolysis In Myocardial Infarction risk index predicted in-hospital mortality in non–ST-segment elevation myocardial infarction, 52
Myocardial revascularization
Early revascularization improved long-term survival after myocardial infarction with cardiogenic shock, 59

N

Nursing homes
A nurse-led clinical pathway reduced hospitalizations in nursing home residents with pneumonia, 81
Nutrition
Dietetic assistants improved postoperative clinical outcomes in older women with hip fracture, 40

O

Obesity
Laparoscopic gastric band surgery was more effective than an intensive nonsurgical intervention for weight loss in mild-to-moderate obesity, 41
Orthomyxoviridae infections
Review: Antiviral agents reduce risk for influenza in healthy adults and alleviate symptoms faster than placebo, 45
Osteoarthritis, knee
Glucosamine and chondroitin sulfate did not improve pain in osteoarthritis of the knee, 17
Osteoporosis
Review: Human parathyroid hormone reduces fractures and increases bone mineral density in severe osteoporosis, 71
Otitis media, suppurative
Review: Topical quinolones are better than systemic antibiotics for chronic suppurative otitis media at up to 2 weeks, 48

P

Pain
Glucosamine and chondroitin sulfate did not improve pain in osteoarthritis of the knee, 17
Review: Opioids are more effective than placebo but not other analgesics for chronic noncancer pain, 74
Parathyroid hormone
Review: Human parathyroid hormone reduces fractures and increases bone mineral density in severe osteoporosis, 71
Paroxetine
Maintenance treatment with paroxetine, but not psychotherapy, prevented recurrent major depression in older persons, 13
Patient compliance
Review: Good adherence (compared with poor adherence) to drug therapy is associated with a reduction in mortality, 80

Patient education
Yoga improved function and reduced symptoms of chronic low-back pain more than a self-care book, 16
Patient readmission
1-year mortality after first hospitalization for heart failure was similar in patients with preserved or reduced ejection fraction, 78
Phytotherapy
Review: Horse chestnut seed extract is effective for symptoms of chronic venous insufficiency, 20
Plant extracts
Saw palmetto did not differ from placebo for benign prostatic hyperplasia in men, 12
Platelet aggregation inhibitors
Abciximab reduced death, MI, and urgent target vessel revascularization in non–ST-segment elevation acute coronary syndromes, 8
Aspirin plus dipyridamole was more effective than aspirin alone for preventing vascular events after minor cerebral ischemia, 57
Clopidogrel plus aspirin did not differ from aspirin alone for reducing MI, stroke, and CV death in high-risk atherothrombosis, 33
Clopidogrel plus aspirin was inferior to oral anticoagulation for preventing vascular events in atrial fibrillation, 58
Pneumonia
A nurse-led clinical pathway reduced hospitalizations in nursing home residents with pneumonia, 81
Oral decontamination with chlorhexidine reduced ventilator-associated pneumonia in high-risk patients, 68
Polysaccharides
Fondaparinux reduced death or reinfarction in acute ST-segment elevation myocardial infarction, 29
Fondaparinux was noninferior to enoxaparin for death, MI, and refractory ischemia but reduced bleeding in angina and non-STEMI, 31
Postoperative complications
Dietetic assistants improved postoperative clinical outcomes in older women with hip fracture, 40
Review: Lung expansion modalities and selective nasogastric decompression prevent postoperative pulmonary complications, 38
Review: Surgical site, advanced age, and comorbid conditions increase risk for postoperative pulmonary complications, 37
Posture
Review: Sparse evidence supports lifestyle modifications for reducing symptoms of gastroesophageal reflux disease, 44
Probiotics
Review: Probiotics are effective for prevention of antibiotic-associated diarrhea and treatment of *Clostridium difficile* disease, 46
Prostatic hyperplasia
Saw palmetto did not differ from placebo for benign prostatic hyperplasia in men, 12
Psychotherapy
Maintenance treatment with paroxetine, but not psychotherapy, prevented recurrent major depression in older persons, 13
Pulmonary disease, chronic obstructive
Formoterol in addition to tiotropium improved airflow obstruction in chronic obstructive pulmonary disease, 39
Pulmonary embolism
A clinical prediction model was sensitive but not specific for predicting mortality in pulmonary embolism, 53
Multidetector CTA with venography was more sensitive for diagnosing pulmonary embolism than CTA alone, 76
The revised Geneva score predicted pulmonary embolism in patients with shortness of breath or chest pain,025
Pyrroles
Atorvastatin did not prevent cardiovascular events in type 2 diabetes, 62

R

Raloxifene
Raloxifene and tamoxifen had similar efficacy for preventing invasive breast cancer in women at increased risk, 72
Raloxifene produced both harms and benefits in postmenopausal women, with no reduction in cardiovascular disease risk, 73
Renal insufficiency, chronic
Benazepril was effective and safe for advanced chronic kidney disease without diabetes, 19
Renin-angiotensin system
Review: The renoprotective effects of ACE inhibitors and ARBs independent of blood pressure control are uncertain, 36
Respiratory distress syndrome, adult
A conservative fluid management strategy did not affect risk for death but shortened duration of ventilation in acute lung injury, 69
Pulmonary artery catheters and central venous catheters did not differ for mortality in acute lung injury, 70
Respiratory insufficiency
Review: Lung expansion modalities and selective nasogastric decompression prevent postoperative pulmonary complications, 38
Review: Surgical site, advanced age, and comorbid conditions increase risk for postoperative pulmonary complications, 37
Ribavirin
Review: Ribavirin is not better than placebo in chronic hepatitis C infection, 47

S

Sciatica
Lumbar epidural corticosteroid injections provided only short-term relief of symptoms and pain in unilateral sciatica, 15
Scopolamine derivatives
Formoterol in addition to tiotropium improved airflow obstruction in chronic obstructive pulmonary disease, 39
Seeds
Review: Horse chestnut seed extract is effective for symptoms of chronic venous insufficiency, 20
Selective estrogen receptor modulators
Raloxifene and tamoxifen had similar efficacy for preventing invasive breast cancer in women at increased risk, 72
Raloxifene produced both harms and benefits in postmenopausal women, with no reduction in cardiovascular disease risk, 73
Self administration
Review: Self-monitoring increases the efficacy and safety of anticoagulant therapy, 1
Self care
Yoga improved function and reduced symptoms of chronic low-back pain more than a self-care book, 16
Serotonin uptake inhibitors
Maintenance treatment with paroxetine, but not psychotherapy, prevented recurrent major depression in older persons, 13
Shock, cardiogenic
Early revascularization improved long-term survival after myocardial infarction with cardiogenic shock, 59
Sleep apnea, obstructive
Review: Oral appliances are better than control appliances, but not CPAP, in the obstructive sleep apnea–hypopnea syndrome, 43
Sleep initiation and maintenance disorders
Review: Sedative–hypnotics increase adverse effects more than they improve sleep quality in older persons with insomnia, 14

Stomatitis
Review: Prophylactic interventions reduce oral mucositis in patients with cancer receiving chemotherapy or radiation therapy, 75
Stroke volume
1-year mortality after first hospitalization for heart failure was similar in patients with preserved or reduced ejection fraction, 78
Sulfonylurea compounds
Adjunctive inhaled insulin before meals improved glycemic control more than adjunctive metformin in type 2 diabetes mellitus, 67

T

Tamoxifen
Disease-free survival was greater with letrozole than tamoxifen in postmenopausal women with early breast cancer, 11
Raloxifene and tamoxifen had similar efficacy for preventing invasive breast cancer in women at increased risk, 72
Tetrazoles
Clopidogrel plus aspirin was inferior to oral anticoagulation for preventing vascular events in atrial fibrillation, 58
Thrombin
Review: Low-molecular-weight heparin is more effective than unfractionated heparin for preventing reinfarction in STEMI, 9
Thrombosis
Clopidogrel plus aspirin did not differ from aspirin alone for reducing MI, stroke, and CV death in high-risk atherothrombosis, 33
Review: Low-molecular-weight heparin is more effective than unfractionated heparin for preventing reinfarction in STEMI, 9
Review: Selective COX-2 inhibitors increase vascular events more than placebo and naproxen, but not more than other NSAIDs, 66
Ticlopidine
Abciximab reduced death, MI, and urgent target vessel revascularization in non–ST-segment elevation acute coronary syndromes, 8
Clopidogrel plus aspirin did not differ from aspirin alone for reducing MI, stroke, and CV death in high-risk atherothrombosis, 33
Clopidogrel plus aspirin was inferior to oral anticoagulation for preventing vascular events in atrial fibrillation, 58
Tomography, spiral computed
Review: MRI is more sensitive but less specific than ultrasonography or spiral CT for diagnosis of hepatocellular carcinoma, 21
Tomography, x-ray computed
Multidetector CTA with venography was more sensitive for diagnosing pulmonary embolism than CTA alone, 76
Triamcinolone acetonide
Lumbar epidural corticosteroid injections provided only short-term relief of symptoms and pain in unilateral sciatica, 15
Triazoles
Disease-free survival was greater with letrozole than tamoxifen in postmenopausal women with early breast cancer, 11
Tumor markers, biological
Review: MRI is more sensitive but less specific than ultrasonography or spiral CT for diagnosis of hepatocellular carcinoma, 21
Tumor necrosis factor-alpha
Review: Anti–tumor necrosis factor antibody therapy for rheumatoid arthritis increases risk for serious infection and malignancy, 65
Tympanic membrane perforation
Review: Topical quinolones are better than systemic antibiotics for chronic suppurative otitis media at up to 2 weeks, 48

U

Ultrasonography
Review: MRI is more sensitive but less specific than ultrasonography or spiral CT for diagnosis of hepatocellular carcinoma, 21

V

Vascular diseases
Lowering homocysteine with folic acid and B vitamins did not prevent vascular events in vascular disease, 2
Venous insufficiency
Review: Horse chestnut seed extract is effective for symptoms of chronic venous insufficiency, 20
Venous thrombosis
Multidetector CTA with venography was more sensitive for diagnosing pulmonary embolism than CTA alone, 76
Review: The Wells clinical prediction guide and D-dimer testing predict deep venous thrombosis, 24
Ventilators, mechanical
Oral decontamination with chlorhexidine reduced ventilator-associated pneumonia in high-risk patients, 68
Ventricular dysfunction, left
Review: Electrocardiography, BNP, and N terminal-pro BNP are more sensitive than specific for chronic left ventricular systolic dysfunction, 22
Vitamins
Calcium plus vitamin D did not prevent hip fracture or colorectal cancer in postmenopausal women, 4
Lowering homocysteine with folic acid and B vitamins did not prevent vascular events in vascular disease, 2
Lowering homocysteine with folic acid and B vitamins did not prevent vascular events after myocardial infarction, 3

W

Warfarin
Review: Self-monitoring increases the efficacy and safety of anticoagulant therapy, 1
Weight loss
Review: Sparse evidence supports lifestyle modifications for reducing symptoms of gastroesophageal reflux disease, 44

Y

Yoga
Yoga improved function and reduced symptoms of chronic low-back pain more than a self-care book, 16